

MOVIPREP Laxative instructions (afternoon appointment)

Information and advice for patients preparing for a colonoscopy or sigmoidoscopy

Bowel Cancer Screening Programme

These are instructions telling you how to take MOVIPREP in preparation for your colonoscopy or sigmoidoscopy.

You are also advised to read any patient information leaflet provided by the manufacturer.

What is MOVIPREP?

MOVIPREP is a strong laxative that you need to take before your colonoscopy or sigmoidoscopy. It is powder which you must mix with water and drink. We will have sent you two clear sealed plastic bags containing the MOVIPREP with your appointment letter or given to you at an appointment. MOVIPREP will give you diarrhoea so you should stay close to a toilet once you have taken it.

What are the benefits of MOVIPREP?

Taking MOVIPREP will empty your bowel so that the lining can be examined properly during your colonoscopy or sigmoidoscopy.

What are the risks of taking MOVIPREP?

The risk of taking MOVIPREP is that it can cause side effects such as nausea (feeling sick), vomiting, diarrhoea, bloating, abdominal (tummy) pain, sleep disturbance and anal irritation. Applying a barrier cream such as zinc and castor oil, or using wet wipes after each bowel movement, can help prevent soreness of the bottom.

Vomiting and diarrhoea can lead to fluid loss (dehydration) with dizziness, headache and confusion if you do not have enough fluid and salt to replace what has been lost.

If you feel unwell while taking MOVIPREP please contact your GP. If your symptoms are severe, please go to A&E or call 999 in an emergency.

What are the risks of not taking MOVIPREP?

If you do not take MOVIPREP correctly, your bowel will not be cleansed enough to ensure a safe and effective examination, which may mean we need to cancel your procedure.

Are there any alternatives to taking MOVIPREP?

There are alternative types of laxative available but these have the same effect. MOVIPREP is the one we recommend. If you are concerned about taking MOVIPREP you can contact the Bowel Cancer Screening team.

MOVIPREP Laxative instructions (afternoon appointment)

Information and advice for patients preparing for a colonoscopy or sigmoidoscopy

Bowel Cancer Screening Programme

Before taking MOVIPREP

Please let the doctors or nurses know about the medication that you are taking before you take MOVIPREP.

If you take diabetic tables or insulin and have not received a diabetic leaflet or you are on tablets that thin the blood such as warfarin, rivaroxaban, dabigatran, clopidogrel/Plavix, prasugrel, tricagrelor or apixaban, please contact the bowel cancer screening team before you take the MOVIPREP.

Medications

Iron tablets last dose on

Anticoagulant last dose on

Antiplatelet last dose on

Diabetic care - leaflet given Yes / No

Other stop on

Other start on

MOVIPREP Laxative instructions (afternoon appointment)

Information and advice for patients preparing for a colonoscopy or sigmoidoscopy

Bowel Cancer Screening Programme

Three days before the test

Date

You need to follow a low residue diet for at least three days before the test to help your bowel empty properly for the colonoscopy or sigmoidoscopy. A low residue diet is made up of foods that are easy to digest and avoids foods that are high in fibre.

High Fibre foods you CAN'T eat	Low residue foods you CAN eat
<ul style="list-style-type: none"> • Bread – wholemeal, high bran, granary, high fibre or half wholegrain • Wholemeal pittas, chapattis, crumpets, scones and muffins • Cereal bars, digestive biscuits, fruit cake, fig rolls • Wholemeal lasagne/pasta • Brown rice • High fibre or bran crisp breads and crackers • Wholemeal, wholegrain bran or high fibre cereals, porridge and muesli • Nuts • Fruit – fresh, frozen, cooked and dried • Beans, lentils and chick peas • Potatoes with skin on 	<ul style="list-style-type: none"> • White cereals such as cornflakes, frosties or rice based cereals • White rice, white pasta or white noodles • Plain white biscuits such as rich tea, custard creams, nice or malted milk • White bread, crumpets and scones (without dried fruit) • Well cooked meat, fish, cheese, egg or tofu • Potatoes without the skin • Butter or margarine • Ice cream, custard, rice pudding or clear jelly

The day before the test

Date

Do not eat anything after 1 pm (after lunch), even if you feel hungry, because your bowel needs to be completely empty for you examination. You can have clear soups, clear soft drinks, clear cordials, black tea, herbal tea or coffee without milk. Sugar can be added if you like but all drinks must not contain fibre.

These drinks can be taken until 2 hours before your procedure.

7 pm

At about 7 pm, make up your first litre of MOVIPREP (sachets A and B) and drink it over 1-2 hours

It is important to drink an additional 500ml of water or clear fluid with each litre of MOVIPREP during the evening.

Allow two hours for MOVIPREP to work after finishing the first litre, before going to bed. You can go to bed when you stop going to the toilet (for example 10 pm).

If you do not have a bowel movement within four hours of taking Moviprep, stop the intake and contact the endoscopy unit.

MOVIPREP Laxative instructions (afternoon appointment)

Information and advice for patients preparing for a colonoscopy or sigmoidoscopy

Bowel Cancer Screening Programme

The day of your test

Date

Remember, do not eat anything.


At 6 am make up your second litre of MOVIPREP (sachets A and B) and drink it over 1-2 hours.

It is important to drink an additional 500ml of water or clear fluid with each litre of MOVIPREP.

Once again you will have watery bowel movements which will stop after 1-2 hours, allowing you to leave home in good time for your hospital appointment. Allow two hours for MOVIPREP to work after finishing your second litre, before leaving for your appointment.

How to prepare MOVIPREP

- Open one clear bag and remove Sachets A and B
- Add the contents of BOTH sachet A and B to a measuring container that holds one litre.
- Add water into the container up to the one litre mark and stir until all the powder has dissolved and the MOVIPREP solution is clear or slightly hazy. This may take up to five minutes.
- After Moviprep is dissolved in the water, the solution can be covered and stored in the fridge for 24 hours.
- You may find it easier to drink the solution using a straw.



(Source: Hints and Advice for patients taking MOVIPREP or MOVIPREP Orange, produced by Norgine).

MOVIPREP Laxative instructions (afternoon appointment)

Information and advice for patients preparing for a colonoscopy or sigmoidoscopy

Bowel Cancer Screening Programme

Key points to remember:

- Do not eat anything after lunch on the day before your colonoscopy or sigmoidoscopy until after your hospital appointment;
- Take one litre of MOVIPREP the night before your colonoscopy or sigmoidoscopy;
- Take one litre of MOVIPREP early in the morning of your examination
- To prepare one litre of MOVIPREP pour the contents of one sachet A and one sachet B into a jug, make up to one litre with water (not chilled);
- Drink an additional 500 ml of water or clear fluids with each sachet of MOVIPREP taken;
- If you take any oral medication, do not take it an hour either side of drinking your dose of MOVIPREP.

After you drink MOVIPREP you will have lots of watery bowel movements, so stay near a toilet because you may need to use it urgently.

Always read the label. It is extremely important to complete your bowel preparation fully as instructed because your doctor will need to have a clear view of the inner lining of your bowel.

If you have missed a dose or taken it incorrectly, please phone the Bowel Cancer Screening nurses.

Contact details

Please contact us if you have any questions or concerns.

Sandwell Hospital Bowel Cancer Screening nurses

0121 507 3185/3874

City Hospital Bowel Cancer Screening nurses

0121 507 6002

Queen Elizabeth Hospital Bowel Cancer Screening nurses

0121 371 6993

When out of hours, please contact:

Sandwell Hospital, Priory 5 Gastroenterology Ward 0121 507 3493

City Hospital, Ward D15 Gastroenterology Ward 0121 507 4405

6.30pm-8am Monday-Friday (evening), and weekends.

MOVIPREP Laxative instructions (afternoon appointment)

Information and advice for patients preparing for a colonoscopy or sigmoidoscopy

Bowel Cancer Screening Programme

Sources used for the information in this leaflet:

- Norgine (2020). *Hints & advice on taking Moviprep*. Uxbridge: Norgine Pharmaceuticals limited.
- Norgine (2019). *Information for the user, Moviprep, powder for oral solution*. Uxbridge: Norgine Pharmaceuticals limited.
- Imperial College healthcare NHS trust (2021). *Low residue diet, information for patients, relatives and carers*. London: Imperial College healthcare NHS trust.
- Moviprep (2017). *How to take Moviprep*. Available at: <https://moviprep.salix.com/about-moviprep/taking-moviprep>

(accessed on January 2022).

If you would like to suggest any amendments or improvements to this leaflet please contact the communications department on 0121 507 5303 or email: swb-tr.swbh-gm-patient-information@nhs.net



A Teaching Trust of The University of Birmingham
Incorporating City, Sandwell and Rowley Regis Hospitals
© Sandwell and West Birmingham Hospitals NHS Trust

ML5794
Issue Date: May 2022
Review Date: May 2025