

Having a Flexible Sigmoidoscopy

Information and advice for patients

Bowel Cancer Screening Programme

What is a Flexible Sigmoidoscopy?

A flexible sigmoidoscopy is an examination of the lining of the large bowel (colon). It involves a colonoscope (a thin, flexible tube with a bright light on the end) being passed through your back passage (rectum) and into the left side of your bowel. It allows samples of tissue (a biopsy) to be taken, or the removal of small warty growths (polyps) that may become cancerous at a later date.

You have been advised to have a flexible sigmoidoscopy to help find the cause of your symptoms and as part of the bowel cancer screening programme.

What are the benefits of this procedure?

The benefit of a colonoscopy is, it can help to diagnose the cause of your problems and prevent you from getting bowel cancer in the future, by removing polyps.

What are the risks of the procedure?

It is not unusual to experience bloating and abdominal discomfort during the procedure and for a few hours after.

Occasionally it is not possible to examine all of the bowel required for the procedure and you may require either a repeat procedure or a different test on another day to complete the examination.

There is a small risk that an adverse reaction to the intravenous sedative and pain relief medication can occur if you are having sedation. However, this is a rare complication.

Removing a polyp can sometimes cause bleeding, although this is usually stopped during the procedure. Occasionally bleeding may occur after you have gone home and, even more rarely, could result in heavy bleeding needing a blood transfusion or even less commonly, surgery.

There is a risk of making a small tear (perforation) to your bowel. This is more likely to occur after the removal of a polyp and may require an operation to repair the damage.

Like all tests, there is a risk the procedure will not show up all abnormalities and, on very rare occasions, a significant abnormality may not be identified like a cancer or a polyp.

If you have any questions about the risks of this procedure, please ask your bowel cancer screening nurse.

Having a Flexible Sigmoidoscopy

Information and advice for patients

Bowel Cancer Screening Programme

What are the alternatives to this procedure?

In some cases, depending on your symptoms and condition, a Computerised Tomography (CT) colon scan may be an alternative to having a flexible sigmoidoscopy. However, a CT scan will not allow the direct visual inspection of the lining of the large bowel wall or enable samples to be taken or polyps removed.

What are the risks of not having the procedure?

If you do not undergo this test, we may not be able to detect an abnormality that would require further treatment. Some conditions can only be detected by a camera test rather than CT scans. If you wish to discuss what it would mean to decline this test please speak to your screening nurse.

What do I need to do before I attend for a flexible sigmoidoscopy?

Bowel preparation

To allow a clear view of your bowel it must be empty of faeces (poo). You may have been given bowel preparation to take for this; please follow the instructions carefully and contact your screening nurse if you have any queries. Alternatively, you may be offered an enema at hospital to clear out the lower part of your bowel (rectum).

Medication

If you take diabetic medication (tablets or insulin) and did not understand or receive your diabetic information please contact your screening nurse. If you take warfarin, clopidogrel/plavix, dabigatran, rivaroxiban, prasugrel, ticagrelor, apixoban or any other blood thinner, and did not understand your instructions for stopping medication please contact the screening nurse team.

Please stop taking all iron tablets 7 days before your appointment.

You may continue to take your other usual medication. Please bring any medication you are currently taking (including sprays and inhalers) with you to your appointment, and details of any allergies or reactions to medicines in the past.

What happens when I arrive at the hospital?

Your appointment will last for 2 – 4 hours. When you arrive please report to the reception desk where a receptionist will check your details and direct you to the waiting area. Please do not bring any valuables or jewellery to the hospital. To respect the privacy of other patients, the endoscopy department does not usually allow friends or relatives to stay with you while attending for your test.

Having a Flexible Sigmoidoscopy

Information and advice for patients

Bowel Cancer Screening Programme

Your screening nurse will then explain the procedure to you, to make sure you understand the benefits and possible risks as detailed in this leaflet. The staff will want you to be as relaxed as possible for the test and will not mind answering questions. Provided you are happy for the procedure to be performed, the screening nurse will ask you to sign the consent form to confirm you understand the procedure.

What happens during the test?

The flexible sigmoidoscopy will take place in a private room with only you, the person doing the colonoscopy and nurses present. You will be asked to change into a hospital gown for the test. The test takes about 15 minutes, but may take longer if a polyp is removed.

The procedure is usually without sedation as it is a short test. If you would like to have sedation please inform the nurse staff or the screening nurses when you arrive. If you do have sedation you will not be able to drive yourself home and you will need an adult to stay with you overnight.

The procedure can be done using Entonox (gas and air) or sedation for pain relief and the screening nurse will discuss this with you at your clinic appointment. If you choose to have Entonox you will administer this yourself by sucking through a tube and breathing in and out. Entonox has fewer side effects than sedation and you will recover more quickly. If you choose to have sedation this will be given through a small cannula tube in the back of your hand or in your arm (cannula). The sedation is not an anaesthetic but it is conscious sedation which may make you sleepy and you may not remember the procedure but you will be awake.

You will be given oxygen through small tubes placed gently in your nostrils. A clip will be attached to a finger or ear lobe so that the levels of oxygen in the blood can be monitored.

Your blood pressure may also be measured during the procedure using a small cuff around your arm.

The procedure

1. In the examination room you will be made comfortable on the couch and will be asked to lie on your left side. Your back passage will be examined using a finger before the sigmoidoscope is inserted.
2. When the sigmoidoscope is inserted, air is passed into the bowel to inflate it, which helps to give a clearer view.
3. This may give you a bloated feeling and some tummy pain which should not last too long. You may also feel the sensation of wanting to go to the toilet, but as your bowel

Having a Flexible Sigmoidoscopy

Information and advice for patients

Bowel Cancer Screening Programme

will be empty this is unlikely. You may pass wind and although this may be embarrassing for you, the nursing staff will understand what is causing the wind.

4. You may be asked to change position to your back or other side during the procedure.
5. If a polyp needs to be removed or a biopsy taken, this is performed through the sigmoidoscope and is not painful. The base of the polyp is usually cauterised (burnt) to reduce the risk of bleeding.

What happens after the test?

After the test you will need to stay in the endoscopy unit to rest for up to 1 hour.

Going home

Before you go home, the wind pains and bloating should have settled. It is important that you tell the nurse if they have not, or if they are becoming worse. If you have had sedation it is essential that a responsible adult comes to pick you up and accompanies you home by car or taxi. Public transport is not suitable.

When you arrive home it is important to rest quietly for the remainder of the day with someone to look after you overnight. It is advisable you have the following day off work. Your screening nurse will contact you by telephone the next working day after the procedure to check you have had no complications.

Sedation can impair your reflexes and judgement. For the first 24 hours following sedation do not:

- Drive a car
- Drink alcohol
- Take sleeping tablets
- Operate any machinery or electrical items; even a kettle
- Work at heights (including climbing ladders or onto chairs)
- Sign any legally binding documents

When will I get the results?

Your screening nurse will explain the test results to you before you are discharged. If a biopsy or polyp has been removed, this will be sent to the laboratory for testing and you will be told the results of this within 7 days. You will also be given a written report of the procedure and instructions as to what to do if you have any problems following the test. If you need one, you will be given a clinic appointment to see the screening nurse to discuss your results.

Having a Flexible Sigmoidoscopy

Information and advice for patients

Bowel Cancer Screening Programme

How to contact us

Bowel cancer screening nurse

City Hospital - 0121 507 6002

Sandwell Hospital - 0121 507 3185

Queen Elizabeth Hospital - 0121 371 6993

If you are unable to keep your appointment please telephone one of the above numbers as soon as possible so the appointment can be given to another patient.

Further information

For more information about our hospitals and services please see our website

www.swbh.nhs.uk, follow us on Twitter @SWBHnhs and like us on Facebook

www.facebook.com/SWBHnhs and <http://www.facebook.com/swbhbcsp>

Having a Flexible Sigmoidoscopy

Information and advice for patients

Bowel Cancer Screening Programme

Sources used for the information in this leaflet

- Veitch AM, et al. (2021) *Endoscopy in patients on antiplatelet or anticoagulant therapy: British Society of Gastroenterology (BSG) and European Society of Gastrointestinal Endoscopy (ESGE) guideline update*. Gut. 70:1611–1628. doi:10.1136/gutjnl-2021-325184. Available at: https://www.bsg.org.uk/wp-content/uploads/2021/08/BSG_ESGE-antiplatelet-and-anticoagulant-update-2021.pdf (Accessed 15 June 2023).
- GOV.UK (2022) *Bowel cancer screening: having a colonoscopy*. Available at: <https://www.gov.uk/government/publications/bowel-cancer-screening-colonoscopy/bowel-cancer-screening-having-a-colonoscopy-fit#:~:text=Colonoscopy%20can%20find%20bowel%20cancer,may%20take%20around%202%20hours>. (Accessed 15 June 2023)
- Chilton, A., Rutter, M. (2011). *Quality assurance guidelines for colonoscopy: publication number 6*. [Online]. Sheffield: NHS cancer screening programmes. Available at: <https://www.gov.uk/government/publications/bowel-cancer-screening-colonoscopy-quality-assurance> (Accessed 30 June 2023).

If you would like to suggest any amendments or improvements to this leaflet please contact SWB Library Services on ext 3587 or email swbh.library@nhs.net.



A Teaching Trust of The University of Birmingham
Incorporating City, Sandwell and Rowley Regis Hospitals
© Sandwell and West Birmingham NHS Trust

ML6090

Issue Date: August 2023

Review Date: August 2026